

Northwest Urology Clinic Inc., PS
1311 East Division Street * Mount Vernon, WA 98274
1213-24th Street, Suite #600 * Anacortes, WA 98221
MVO TEL 360-424-7991 ANA TEL 360-293-2173
MVO FAX 360-428-4377 ANA TEL 360-293-0910

BCG BLADDER INSTILLATION

TIME TO REPORT...DAY _____ DATE _____ TIME _____ AM/PM

TIME TO REPORT...DAY _____ DATE _____ TIME _____ AM/PM

TIME TO REPORT...DAY _____ DATE _____ TIME _____ AM/PM

TIME TO REPORT...DAY _____ DATE _____ TIME _____ AM/PM

TIME TO REPORT...DAY _____ DATE _____ TIME _____ AM/PM

TIME TO REPORT...DAY _____ DATE _____ TIME _____ AM/PM

SCHEDULED AT MT. VERNON ANACORTES STANWOOD

To make sure each treatment has the best chance to work, you will be asked to keep the BCG solution in your bladder for 2 hours after receiving your therapy.

POST INSTILLATION INSTRUCTIONS:

Please follow these directions for 6 hours after each BCG treatment:

1. Sit on the toilet seat to urinate.
2. Immediately after urinating, add 2 cups of undiluted chlorine bleach to the toilet and close the lid.
3. Let the bleach stand for 15 minutes before flushing to disinfect the toilet.
4. Drink plenty of water to flush any remaining BCG bacteria out of your bladder
5. Disinfect the toilet with 2 cups of undiluted chlorine bleach for 15 minutes each time you urinate for the 6 hours after treatment.

SIDE EFFECTS TO BCG:

Tell your doctor if you experience the following:

- Difficulty urinating, frequent urination, blood in your urine.
- Chills, night sweats, severe sluggishness.
- Persistent fever for more than 2 days.
- Joint pain, cough, skin rash.

Darrell Cornelius, M.D.

Winston Jones, M.D.

Terry Milam, M.D.

Macintosh HD:Users:Art:Documents:Microsoft User Data:Saved Attachments:BCG
TREATMENT INSTRUCTIONS.doc